ISSN (print) 0093-4666 © 2015. Mycotaxon, Ltd. ISSN (online) 2154-8889

MYCOTAXON

http://dx.doi.org/10.5248/130.237

Volume 130, pp. 237–239 *January–March 2015*

Phyllachora hainanensis sp. nov. from China

Na Liu[1]*, Le Wang[1], Gu Huang[1,2], & Lin Guo[2]

[1] *College of Biological Engineering, Henan University of Technology, Lianhua Street, High & New Tech Development District, Zhengzhou 450001, China*

[2] *State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences, No.1 Beichen West Road, Chaoyang District, Beijing 100101, China*

* Correspondence to: *liuna3456@163.com*

Abstract — A new species, *Phyllachora hainanensis* on *Phyllanthus emblica* (*Phyllanthaceae*), causing tar spots is reported from Hainan Province, China. *Phyllachora hainanensis* differs from other similar species in the size and shape of ascospores.

Key words — *Sordariomycetes*, taxonomy

A new species of *Phyllachora* (*Sordariomycetes*) on leaves of *Phyllanthus emblica* was collected from Hainan Province, China, in November 2007. Worldwide, only *P. phyllanthophila* and *P. phyllanthophila* var. *egregia* have previously been reported from the host genus *Phyllanthus* (Saccardo & Saccardo 1905: 834, Saccardo & Trotter 1913: 419).

Phyllachora hainanensis Na Liu & G. Huang, **sp. nov.** Plate 1

Fungal Name FN570110

Differs from *Phyllachora phyllanthophila* by its larger ascospores.

Type: China, Hainan Province, Xiawangling, alt. 800 m, on leaves of *Phyllanthus emblica* L. (*Phyllanthaceae*), 13 XI 2007, Shuanghui He, L. Guo & Z.Y. Li 2013 (HMAS 245338, **holotype**).

Etymology: From Hainan Province, where the new species was collected.

Ascomata epigenous, raised, shiny, black, sparse, subglobose, ellipsoidal or irregular, approx. 370 × 150–260 µm diam., immersed in the mesophyll, usually with a single locule, ostiole conspicuous with a conical neck extending through the host epidermis and cuticle, without periphyses. Asci arise from the basal and lateral walls of the ascoma, upper wall to 50 µm thick, composed of epidermal cells occluded by melanised material. Asci 60–100 × 30–70 µm,



Plate 1. *Phyllachora hainanensis* (HMAS 245338, holotype).
1. Ascomata; 2–4. Asci and ascospores.

8-spored, clavate, obtuse at apex, long pedunculate, thin-walled at maturity, unitunicate. Ascospores biseriate or irregularly biseriate, 11–18 × 9–13 µm, ellipsoidal or subglobose, one-celled, hyaline, thin-walled, smooth, without a gelatinous sheath. Anamorph not seen.

Comments—Subglobose ascospores are rare in *Phyllachora*. Only one other species, *P. yunnanensis* Na Liu & L. Guo, collected on *Lespedeza* (*Fabaceae*) from Yunnan Province of China, has been reported as having subglobose (sometimes ellipsoidal) ascospores. *Phyllachora yunnanensis* differs from *P. hainanensis* in its smaller (7–13.5 × 5–9 µm) ascospores (Liu & Guo 2006).

Phyllachora hainanensis is the first *Phyllachora* species in China discovered on a host species in the *Phyllanthaceae*. Most *Phyllachora* species reported

in China occur on hosts in the *Poaceae, Fabaceae,* and *Cyperaceae* (Tai 1979, Teng 1963, Lan et al. 2012, Cui 2009). The only other reports of *Phyllachora* on *Phyllanthus* are *Phyllachora phyllanthophila* Henn. on *Phyllanthus* sp. from Peru (Hennings 1904: 250) and *P. phyllanthophila* var. *egregia* Rehm on *Phyllanthus* sp. from Brazil (Rehm 1907: 530). These taxa were subsequently transferred to *Catacauma* as *C. phyllanthophilum* (Henn.) Theiss. & Syd. and *C. egregium* (Rehm) Theiss. & Syd. (Theissen & Sydow 1915: 390–391), but *Catacauma* is now synonymised with *Phyllachora* (von Arx & Müller 1954: 212) and *P. phyllanthophila* var. *egregia* is synonymised with *P. phyllanthophila* (Index Fungorum 2015). The publications cited above provide different sets of measurements that can be summarised as 35–60 × 12–20 µm for asci, and 10–14 × 5–9 µm for ascospores; thus *P. phyllanthophila* differs from *P. hainanensis* by its smaller asci and its narrower, ellipsoidal to subovoid ascospores.

Acknowledgments

The authors would like to thank Dr. Roger Shivas (Brisbane, Australia) and Guangyu Sun (Northwest Agriculture and Forestry University) for serving as pre-submission reviewers, Drs. Shuanghui He and Zhenying Li for collecting the specimen, Dr. Shaun Pennycook (Auckland, New Zealand) for nomenclatural review, and Mr. Ziyu Cao (Institute of Botany, Chinese Academy of Sciences) for identifying the host plants. This study was supported by the National Natural Science Foundation of China (No. 31300015) and Natural Science Research Project of Henan Province (No. 132300410350).

Literature cited

von Arx JA, Müller E. 1951. Die Gattungen der amerosporen Pyrenomyceten. Beiträge zur Kryptogamenflora der Schweiz 11(1). 434 p.

Cui JQ. 2009. Taxonomy and molecular phylogenetics of *Colletotrichum* and *Phyllachora*. Thesis for Master Degree of Northwest Agriculture and Forestry University. 66 p.

Hennings P. 1904. Fungi amazonici II. a cl. Ernesto Ule collecti. Hedwigia 43: 242–273.

Index Fungorum. 2015. http://www.indexfungorum.org/names/Names.asp [accessed: February 2015].

Lan JQ, Lu HJ, Wu J., Li XL, Zhang T, Kong CC, Kang ZS. 2012. Three new records of *Phyllachora* in China. Mycosystema 31: 639–641. http://dx.doi.org/10.13346/j.mycosystema.2012.04.021

Liu N, Guo L. 2006. A new species of *Phyllachora* (*Phyllachorales*) on *Leguminosae* from China. Mycotaxon 97: 111–114.

Rehm H. 1907. Ascomycetes novi. Annales Mycologici 5: 516–546.

Saccardo PA, Saccardo D. 1905. Supplementum universale, pars VI. Hymenomycetae – Laboulbeniomycetae. Sylloge Fungorum 17. 991 p.

Saccardo PA, Trotter A. 1913. Supplementum universale, pars IX. Ascomycetae – Deuteromycetae. Sylloge Fungorum 22. 1612 p.

Tai FL. 1979. Sylloge Fungorum Sinicorum. Science Press, Beijing, 1527 p.

Teng SC. 1963. Fungi in China. Science Press, Beijing. 808 p.

Theissen F, Sydow H. 1915. Die *Dothideales.* Kritisch-systematische Originaluntersuchungen. Annales Mycologici 13: 149–746.